

# The New Way to Define Key Oncogenic Drivers of Small Cell Lung Cancer

<sup>†</sup>Kee-Beom Kim

BK21 FOUR KNU Creative BioResearch Group, School of Life Sciences, Kyungpook National University, Daegu 41566, Korea



Received: November 24, 2022

Revised: January 18, 2023

Accepted: February 14, 2023

## <sup>†</sup>Corresponding author

Kee-Beom Kim

BK21 FOUR KNU Creative BioResearch Group, School of Life Sciences, Kyungpook National University, Daegu 41566, Korea  
Tel: +82-53-950-6351  
E-mail: kbk@knu.ac.kr

Copyright © 2023 The Korean Society of Developmental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ORCID

Kee-Beom Kim

<https://orcid.org/0000-0003-0132-8924>

## Conflict of interests

The author declare no potential conflict of interest.

## Acknowledgements

This research was supported by Kyungpook National University Research Fund, 2022.

## Authors' contributions

The article is prepared by a single author.

## Ethics approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## Abstract

Small-cell lung cancer (SCLC) continues to be the deadliest of all lung cancer types. Its high mortality is largely attributed to the unchangeable development of resistance to standard chemo/radiotherapies, which have remained invariable for the past 30 years, underlining the need for new therapeutic approaches. Recent studies of SCLC genome revealed a large number of somatic alterations and identified remarkable heterogeneity of the frequent mutations except for the loss of both *RB* and *P53* tumor suppressor genes (TSGs). Identifying the somatic alterations scattered throughout the SCLC genome will help to define the underlying mechanism of the disease and pave the way for the discovery of therapeutic vulnerabilities associated with genomic alterations. The new technique made it possible to determine the underlying mechanism for the discovery of therapeutic targets. To these ends, the techniques have been focused on understanding the molecular determinants of SCLC.

**Keywords:** Small-cell lung cancer (SCLC), pre-cancerous cells (preSC), Somatic engineering mouse model, Clustered regularly interspaced short palindromic repeats (CRISPR)

Lung cancer is the leading cause of cancer deaths among both men and women. The American Cancer Society indicates that there were more than 130,000 deaths from lung cancer in 2022 in the United States alone. Small-cell lung cancer (SCLC) accounts for approximately 10%–15% of all lung cancers and is a high-grade neuroendocrine neoplasm that is distinct from the rare subtypes of large cell neuroendocrine carcinoma and neuroendocrine carcinoid (Lewis et al., 2014; Park et al., 2016; Torre et al., 2016). It is an exceptionally lethal type of lung cancer with a 5-year survival rate of 7%, significantly worse than that of NSCLC (26%) (Byers & Rudin, 2015; Rudin et al., 2021). The high mortality of SCLC patients is mainly due to the highly metastatic nature of the cancer and the invariable resistance to current therapies that have remained largely unchanged for the past 30 years. Despite the recent success of tobacco control policy and smoking cessation campaigns in reducing the disease incidence, the number of the patients will remain high for decades to come, not only because former smokers would still develop the cancer after a long latency but also because approximately 20% of the adult population in the US continues to smoke. While the long tumor latency and the known risk population of SCLC rationalize chemoprevention, the paucity of molecular targets for chemoprevention stems from the poor understanding of the SCLC progression beyond the role of *RB* and *P53* mutations crucial for tumor development. The conventional chemotherapy of cisplatin and etoposide has not significantly improved overall survival in SCLC patients and the development of targeted therapies has been challenging due to the paucity of clinically viable targets. SCLC is mostly driven by loss of tumor


suppressors, including inactivation of RB and P53 and frequent loss of other tumor suppressors such as the RB family members RBL1 and RBL2 (coded by the *RBL1* and *RBL2* and also known as *p107* and *p130*, respectively) (Peifer et al., 2012; Rudin et al., 2012; George et al., 2015; Augert et al., 2017). A better way to identify the mechanism underlying the development of resistance may lead to the development of a novel therapeutic strategy. In addition, an advanced understanding of the mechanisms of tumor initiation and early-stage progression may attribute to the development of novel means for early detection and prevention.

Functional characterization of the relevant cancer genes has been of prime interest but challenging due to lack of systematic and informed ways of defining oncogenic drivers. Particularly, defining gene function during early cancer progression is extremely difficult in most cancer types, because pre-cancerous lesion is often not accessible for functional study and cell lines derived from full-blown tumors may not be an ideal model (Weinstein et al., 2013). Genetically-engineered mouse model (GEMM) has provided a unique opportunity to understand the mechanisms of cancer progression and identify potential therapeutic targets (Friedel et al., 2007; Hansen et al., 2008; Walrath et al., 2010). However, the GEMM usually requires combining multiple alleles through a new gene, limiting characterization of a large number of mutations and also interactions among heterogeneous genetic factors driving tumor development. Recent studies present a 2-step approach for testing the disease relevance of uncharacterized somatic alterations found in SCLC. First, in an *in vitro* model, we induce candidate mutations in pre-cancerous cells (preSC) derived from the GEMM and characterize cellular phenotypes for transformation (Kim et al., 2016; Jia et al., 2018; Kim et al., 2020; Kim et al., 2022a). Second, we suggest in an *in vivo* model where we induce the *in vitro* prioritized mutations in the GEMM, using adenoviral delivery of clustered regularly interspaced short palindromic repeats (CRISPR)-Cre hybrid system (Kim et al., 2022b). The successful outcome of the models will lead to identification of key drivers of SCLC, paving the way for future experiments aimed at novel strategies for prevention and treatment, and also introduce new approaches that can be readily applied to study a wide range of cancer types that face similar research challenge.

## ISOLATION OF PRE-CANCEROUS PRECURSOR OF SMALL-CELL LUNG CANCER (SCLC)

GEMM was developed based on the frequent RB and P53 mutations ( $\geq 90\%$ ) in human SCLC (Peifer et al., 2012; Rudin et al., 2012; George et al., 2015; Augert et al., 2017). Infection of lung epithelial cells of mice carrying floxed alleles of *Rb* and *p53* (*Rb*<sup>lox/lox</sup>; *p53*<sup>lox/lox</sup>) with adenovirus expressing Cre-recombinase (Ad-Cre) results in loss of function, and the 'Rb/p53' mutant lung develops tumors that closely resemble histopathological features similar to those of human SCLC, including neuroendocrine differentiation, genome-wide gene expression, and metastasis patterns. Recently, we developed a variant of the GEMM bearing additional loss of *Rbl2* (a Rb family member, also known as *p130*). This 'Rb/p53/Rbl2' model develops more SCLC tumors with shorter latency compared to the 'Rb/p53' model while the two models are almost identical at the molecular level and both resembles human SCLC (Schaffer et al., 2010). Recent validation of lung neuroendocrine cells (LNECs) as SCLC cell-of-origin enables the characterization of mutant cells at any stage of tumor progression using the mouse model combined with neuroendocrine-lineage specific tools (Kim et al., 2016). Using a neuroendocrine-specific reporter mouse strain (Chga-GFP) that expresses GFP under control of the transcription regulatory elements of *Chromogranin A* gene (*Chga*), a pan-neuroendocrine marker (Oberge et al., 2015; Balanis et al., 2019) specifically labeled LNECs (Fig. 1). To facilitate characterization of mutant cells at any stage of tumor progression,

we crossed the Chga-GFP strain with the GEMM of SCLC to generate ChgA-GFP/Rb/p53/Rbl2 and isolated GFP-positive cells from the lungs one month after Ad-Cre infection, when the lungs did not show macroscopic lesions, or full-blown tumors. Notably, the cells from early stage grew in monolayers, whereas the tumor cells isolated from fully grown tumors formed spheres or aggregates and grew in suspension or loosely attached to culture dish, similarly to human SCLC cells. These results indicate that these cells are immortalized due to the deletions of *Rb* and *p53* (a common cause of cell immortalization), yet not tumorigenic because these cells, unlike SCLC cells, lack oncogenic factors, such as *L-Myc*, but maintain a high level of *E-cadherin*, while both cell types express Chga (Cardnell et al., 2017; Mollaoglu et al., 2017; Chalishazar et al., 2019; Gay et al., 2021). Therefore, we postulate that these cells are pre-cancerous precursors of SCLC tumor (herein 'preSC'). This model was developed based on the pre-malignancy of preSC derived from the GEMM of SCLC using a lineage-labeling tool. It serves as an efficient system to characterize candidate drivers for their abilities to transform preSC to tumorigenic cells in culture and allograft experiment with immune competent mice. In addition, comparative gene expression profiling of preSC and tumor cells, so called "apple-to-apple approach", enables identification of genes and pathways that are directly related to the pathogenesis of the cancer. Given a turnaround time of 2 months on average to test a gene as well as the capacity of simultaneously testing multiple genes, this approach should enable functional interrogation of a large number of genes, increasing the chance of finding key oncogenic drivers.




**Fig. 1. In vitro prioritization of oncogenic mutations of human SCLC.** Toward understanding a complex network of genes and pathways controlling SCLC progression, one of our initial goals is to systematically prioritize 'oncogenic' driver mutations with the eventual goal of functional validation *in vivo*. The system will use CRISPR-technology in the preSC cells to define candidate TSGs that were recently identified in the analyses of SCLC genome but remain uncharacterized. Results of the proposed experiments will identify novel genes that may be critical for the development of SCLC. CRISPR, clustered regularly interspaced short palindromic repeats; RFP, red fluorescent protein; SCLC, small-cell lung cancer; TSG, tumor suppressor gene.

## IN VIVO VALIDATION SMALL-CELL LUNG CANCER (SCLC) MUTATIONS USING AN ADENOVIRAL CLUSTERED REGULARLY INTERSPACED SHORT PALINDROMIC REPEATS (CRISPR)-CRE HYBRID SYSTEM

CRISPR, a microbial nuclease system involved in defense against invading phages, has been modified as a molecular tool for genome engineering at the cellular or organic level, simply using guide RNA and CRISPR-associated nuclease (Cas) (Cong et al., 2013; Mali et al., 2013). Briefly, a guide RNA with a complementary sequence recruits Cas to a target genomic sequence. Cas then causes a double-stranded break that triggers the error-prone repair mechanisms of non-homologous end-joining. The repair process often results in frame shift, deletion, or insertion mutations.

Unlike modeling gene amplification via lentiviral transduction, recapitulation of somatic mutation *in vivo* has difficult, which limits characterization of tumor suppressor genes (TSGs). Here we propose to develop a strategy for efficient functional validation of oncogenic drivers *in vivo*, by introducing candidate mutations in the GEMM using CRISPR-mediated genome editing, and characterize tumors whether they resemble autochthonous tumors of the mouse model as well as human tumors. We engineered a novel CRISPR-Cre hybrid vector by genetically fusing the CRISPR cassette to Cre through a self-cleaving peptide T2A derived from *Thosea asigna virus* (Fig. 2) (Kim et al., 2022b). The vector expresses guide RNA and Cas-2A-Cre fusion protein in the cells. Self-cleavage of 2A separate Cas and Cre that result in CRISPR-mediated mutation and deletion of floxed *Rb* and *p53* alleles, respectively, in the same cell. For *in vivo* use, we incorporated



**Fig. 2.** *In vivo* validation of SCLC mutations using an adenoviral CRISPR-Cre hybrid system. The system proposes to develop a strategy for efficient validation of oncogenic drivers *in vivo*, by introducing candidate mutations in the GEMM using CRISPR-mediated genome editing. Functional validation of candidate TSGs *in vivo* will set the stage for a comprehensive understanding of SCLC progression. TSG, tumor suppressor gene; CRISPR, clustered regularly interspaced short palindromic repeats; SCLC, small-cell lung cancer; GEMM, genetically-engineered mouse model.

this hybrid system in an adenoviral vector (pAd/PL-DEST) using Gateway cloning kit from Life Technologies (Russell et al., 2000). We validated the adenoviral vector using a mouse Cre-reporter cell line that expresses YFP upon Cre-mediated deletion of lox-stop-lox cassette that usually blocks the gene expression. We confirmed the mutations in the target region of the genome of the infected cells while Ad-Cre control did not induce mutation (Kim et al., 2022b). This adenoviral vector will produce the serotype 5 virus (Ad5), the same type as Ad-Cre that we routinely use to infect the lungs of GEMM. High-titer virus particles are produced and accurately titrated by Vector Development Lab at Baylor College of Medicine, the source of Ad-Cre for many lung cancer researches. CRISPR approaches have just merged for the models of liver and non-small cell lung cancers (Xue et al., 2014; Yin et al., 2014; Rogers et al., 2018; Ng et al., 2020). Despite their significance, these approaches still require a combination of a GEMM and a CRISPR system to study many cancer types, including SCLC, that are thought to be initiated by mutation of multiple alleles. Our CRISPR-Cre hybrid system is capable of initiating tumor and altering specific genomic sequence in the same cell. Adenoviral delivery of this system enables functional characterization of candidate gene *in vivo*, overcoming challenges that traditional mouse genetics approaches face to study numerous genetic factors. This combination of the CRISPR-Cre hybrid system with the GEMM is tractable and robust in characterizing a large number of mutations and genes. Our new model and approach will accelerate efforts to understand the biology of SCLC and also other cancer models that are accessible to virus-mediated gene delivery.

## CONCLUSION

We expect increased tumorigenesis in the mice of the experimental group. If the mice develop larger tumors than the controls. This would mean that the introduced mutations caused increased proliferation rate and/or earlier tumor initiation. This can be differentiated by monitoring the mouse lungs using bioimaging instruments over time. In case of accelerated progression, the tumors in the experimental group would display features of higher-grades and be more metastatic compared to those in the controls. If tumor number also increases, that would mean that the genes similar to *E300* normally suppress tumor initiation as well as progression. However, if the tumors of the experimental group is not significantly different from those of controls but have intended mutations, this will indicate that the mutations are not sufficient to drive *in vivo* tumor progression.

Defining patient mutations will advance the field of SCLC, paving the way to mechanistic elucidation and integration of oncogenic pathways and leading to development of a targeted therapy or optimal drug selections for treatment. Furthermore, the new technique, integrating tractable models of tumor progression with high-resolution genomic data, will impact current efforts aimed at discovery of ‘cancer drivers’.

## REFERENCES

- Augert A, Zhang Q, Bates B, Cui M, Wang X, Wildey G, Dowlati A, MacPherson D (2017) Small cell lung cancer exhibits frequent inactivating mutations in the histone methyltransferase KMT2D/MLL2: CALGB 151111 (alliance). *J Thorac Oncol* 12:704-713.
- Balanis NG, Sheu KM, Esedebe FN, Patel SJ, Smith BA, Park JW, Alhani S, Gomperts BN, Huang J, Witte ON, Graeber TG (2019) Pan-cancer convergence to a small-cell neuroendocrine phenotype that shares susceptibilities with hematological malignancies. *Cancer Cell* 36:17-34.E7.
- Byers LA, Rudin CM (2015) Small cell lung cancer: Where do we go from here? *Cancer* 121:664-

672.

- Cardnell RJ, Li L, Sen T, Bara R, Tong P, Fujimoto J, Ireland AS, Guthrie MR, Bheddah S, Banerjee U, Kalu NN, Fan YH, Dylla SJ, Johnson FM, Wistuba II, Oliver TG, Heymach JV, Glisson BS, Wang J, Byers LA (2017) Protein expression of TTF1 and cMYC define distinct molecular subgroups of small cell lung cancer with unique vulnerabilities to aurora kinase inhibition, DLL3 targeting, and other targeted therapies. *Oncotarget* 8:73419-73432.
- Chalishazar MD, Wait SJ, Huang F, Ireland AS, Mukhopadhyay A, Lee Y, Schuman SS, Guthrie MR, Berrett KC, Vahrenkamp JM, Hu Z, Kudla M, Modzelewska K, Wang G, Ingolia NT, Gertz J, Lum DH, Cosulich SC, Bomalaski JS, DeBerardinis RJ, Oliver TG (2019) MYC-driven small-cell lung cancer is metabolically distinct and vulnerable to arginine depletion. *Clin Cancer Res* 25:5107-5121.
- Cong L, Ann Ran F, Cox D, Lin S, Barretto R, Habib N, Hsu PD, Wu X, Jiang W, Marraffini LA, Zhang F (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science* 339:819-823.
- Friedel RH, Seisenberger C, Kaloff C, Wurst W (2007) EUCOMM—The European conditional mouse mutagenesis program. *Brief Funct Genomics Proteomics* 6:180-185.
- Gay CM, Allison Stewart C, Park EM, Diao L, Groves SM, Heeke S, et al. (2021) Patterns of transcription factor programs and immune pathway activation define four major subtypes of SCLC with distinct therapeutic vulnerabilities. *Cancer Cell* 39:346-360.E7.
- George J, Lim JS, Jang SJ, Cun Y, Ozretić L, Kong G, et al. (2015) Comprehensive genomic profiles of small cell lung cancer. *Nature* 524:47-53.
- Hansen GM, Markesich DC, Burnett MB, Zhu Q, Dionne KM, Richter LJ, Finnell RH, Sands AT, Zambrowicz BP, Abuin A (2008) Large-scale gene trapping in C57BL/6N mouse embryonic stem cells. *Genome Res* 18:1670-1679.
- Jia D, Augert A, Kim DW, Eastwood E, Wu N, Ibrahim AH, Kim KB, Dunn CT, Pillai SPS, Gazdar AF, Bolouri H, Park KS, MacPherson D (2018) *Crebbp* loss drives small cell lung cancer and increases sensitivity to HDAC inhibition. *Cancer Discov* 8:1422-1437.
- Kim DW, Wu N, Kim YC, Cheng PF, Basom R, Kim D, Dunn CT, Lee AY, Kim K, Lee CS, Singh A, Gazdar AF, Harris CR, Eisenman RN, Park KS, MacPherson D (2016) Genetic requirement for *Myel* and efficacy of RNA Pol I inhibition in mouse models of small cell lung cancer. *Genes Dev* 30:1289-1299.
- Kim KB, Kabra A, Kim DW, Xue Y, Huang Y, Hou PC, Zhou Y, Miranda LJ, Park JI, Shi X, Bender TP, Bushweller JH, Park KS (2022a) KIX domain determines a selective tumor-promoting role for EP300 and its vulnerability in small cell lung cancer. *Sci Adv* 8:eabl4618.
- Kim KB, Kim DW, Kim Y, Tang J, Kirk N, Gan Y, Kim B, Fang B, Park J, Zheng Y, Park KS (2022b) WNT5A–RHOA signaling is a driver of tumorigenesis and represents a therapeutically actionable vulnerability in small cell lung cancer. *Cancer Res* 82:4219-4233.
- Kim KB, Kim Y, Rivard CJ, Kim DW, Park KS (2020) FGFR1 is critical for RBL2 loss-driven tumor development and requires PLCG1 activation for continued growth of small cell lung cancer. *Cancer Res* 80:5051-5062.
- Lally BE, Urbanic JJ, William Blackstock A, Miller AA, Perry MC (2007) Small cell lung cancer: Have we made any progress over the last 25 years? *Oncologist* 12:1096-1104.
- Lewis DR, Check DP, Caporaso NE, Travis WD, Devesa SS (2014) US lung cancer trends by histologic type. *Cancer* 120:2883-2892.
- Mali P, Yang L, Esvelt KM, Aach J, Guell M, DiCarlo JE, Norville JE, Church GM (2013) RNA-guided human genome engineering via Cas9. *Science* 339:823-826.
- Mollaoglu G, Guthrie MR, Böhm S, Brägelmann J, Can I, Ballieu PM, et al. (2017) MYC drives

- progression of small cell lung cancer to a variant neuroendocrine subtype with vulnerability to Aurora kinase inhibition. *Cancer Cell* 31:270-285.
- Ng SR, Rideout WM 3rd, Akama-Garren EH, Bhutkar A, Mercer KL, Schenkel JM, Bronson RT, Jacks T (2020) CRISPR-mediated modeling and functional validation of candidate tumor suppressor genes in small cell lung cancer. *Proc Natl Acad Sci USA* 117:513-521.
- Oberg K, Modlin IM, De Herder W, Pavel M, Klimstra D, Frilling A, Metz DC, Heaney A, Kwekkeboom D, Strosberg J, Meyer T, Moss SF, Washington K, Wolin E, Liu E, Goldenring J (2015) Consensus on biomarkers for neuroendocrine tumour disease. *Lancet Oncol* 16:E435-E446.
- Park JY, Jang SH (2016) Epidemiology of lung cancer in Korea: Recent trends. *Tuberc Respir Dis* 79:58-69.
- Peifer M, Fernández-Cuesta L, Sos ML, George J, Seidel D, Kasper LH, et al. (2012) Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer. *Nat Genet* 44:1104-1110.
- Rogers ZN, McFarland CD, Winters IP, Seoane JA, Brady JJ, Yoon S, Curtis C, Petrov DA, Winslow MM (2018) Mapping the *in vivo* fitness landscape of lung adenocarcinoma tumor suppression in mice. *Nat Genet* 50:483-486.
- Rudin CM, Brambilla E, Faivre-Finn C, Sage J (2021) Small-cell lung cancer. *Nat Rev Dis Primers* 7:3.
- Rudin CM, Durinck S, Stawiski EW, Poirier JT, Modrusan Z, Shames DS, et al. (2012) Comprehensive genomic analysis identifies SOX2 as a frequently amplified gene in small-cell lung cancer. *Nat Genet* 44:1111-1116.
- Russell WC (2000) Update on adenovirus and its vectors. *J Gen Virol* 81:2573-2604.
- Schaffer BE, Park KS, Yiu G, Conklin JF, Lin C, Burkhardt DL, Karnezis AN, Alejandro Sweet-Cordero EA, Sage J (2010) Loss of p130 accelerates tumor development in a mouse model for human small-cell lung carcinoma. *Cancer Res* 70:3877-3883.
- Torre LA, Siegel RL, Jemal A (2016) Lung cancer statistics. In: Ahmad A, Gadgeel S (eds), *Lung Cancer and Personalized Medicine: Current Knowledge and Therapies. Advances in Experimental Medicine and Biology*. Vol. 893. Springer, Cham, pp 1-19.
- Walrath JC, Hawes JJ, Van Dyke T, Reilly KM (2010) Chapter 4. Genetically engineered mouse models in cancer research. *Adv Cancer Res* 106:113-164.
- Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, Shmulevich I, Sander C, Stuart JM (2013) The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet* 45:1113-1120.
- Xue W, Chen S, Yin H, Tammela T, Papagiannakopoulos T, Joshi NS, Cai W, Yang G, Bronson R, Crowley DG, Zhang F, Anderson DG, Sharp PA, Jacks T (2014) CRISPR-mediated direct mutation of cancer genes in the mouse liver. *Nature* 514:380-384.
- Yin H, Xue W, Chen S, Bogorad RL, Benedetti E, Grompe M, Kotliansky V, Sharp PA, Jacks T, Anderson DG (2014) Genome editing with Cas9 in adult mice corrects a disease mutation and phenotype. *Nat Biotechnol* 32:551-553.