

Genetic Differences of Three *Pollicipes mitella* Populations Identified by PCR Analysis

Young-Jae Song and †Jong-Man Yoon

Department of Aquatic Life Medicine, College of Ocean Science and Technology,
Kunsan National University, Gunsan 573-701, Republic of Korea

ABSTRACT : Genomic DNAs were extracted from the turtle leg (*Pollicipes mitella*, 1798) population of Tongyeong, Yeosu and Manjaedo located in the southern sea of Korea. The turtle leg population from Tongyeong (0.929) exhibited higher bandsharing values than did turtle leg from Manjaedo (0.852). The higher fragment sizes (>1,200 bp) are much more observed in the Yeosu population. The number of unique loci to each population and number of shared loci by the three populations, generated by PCR using 7 primers in the turtle leg (*P. mitella*) population of Tongyeong, Yeosu and Manjaedo. Genetic distances among different individuals of the Tongyeong population of the turtle leg (lane 1 –07), Yeosu population of the turtle leg (lane 08–14) and Manjaedo population of the turtle leg (lane 15 –21), respectively, were generated using the CLASSIFICATION option in Systat version 10 according to the bandsharing values and similarity matrix. The dendrogram, obtained by the seven decamer primers, indicated three genetic clusters: cluster 1 (TONGYEONG 01 –TONGYEONG 07), cluster 2 (YEOSU 08–YEOSU 14), and cluster 3 (MANJEDO 15–MANJEDO 21). Tongyeong population could be evidently discriminated with the other two Yeosu and Manjaedo populations among three populations. The longest genetic distance (0.305) was found to exist between individuals' no. 02 of the Tongyeong population and no. 13 of the Yeosu population. It seems to the authors that this is a result of a high degree of inbreeding in narrow region for a long while.

Key words : Genetic clusters, Genetic distance, Inbreeding, Manjaedo, *Pollicipes mitella*, Tongyeong, Turtle leg, Yeosu

INTRODUCTION

The turtle leg (*Pollicipes mitella*, 1798), which belongs to the family Pollicipes, is widely distributed in the rock along the shore areas of the southern sea, Jeju island and Ulleungdo island of Korean Peninsula. Generally, the size, type, stripe pattern, and color of this species vary in accordance with water depth, turbidity, nutrition, growth period, water temperature, and other environmental aspects. Research for artificial production has progressed steadily in many aspects, over-catching, and water pollution

by industries and city sewage. As this species culture industry grows, so doe's interest into the genetics of this species. However, little information currently exists regarding the genetics of this species.

Many molecular/genetic studies have employed this technique, as this PCR method is a reliable, easy, and relatively speedy method for the investigation of numerous genomic DNAs, which can help in the determination of the degree of genetic diversity in a given population. Another benefit of this method is that it requires no prior knowledge of the genome for its efficiency (Welsh

Manuscript received 18 August 2013, Received in revised form 24 August 2013, Accepted 29 August 2013

† Corresponding Author : Jong-Man Yoon, Department of Aquatic Life Medicine, College of Ocean Science and Technology, Kunsan National University, Gunsan 573-701, Republic of Korea. Tel. : +82-63-469-1887, Fax : +82-63-463-9493, E-mail : jmyoon@kunsan.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License(<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

& McClelland, 1990; Welsh et al., 1991; Iyengar et al., 2000; Klinbunga et al., 2000).

In the present study, these authors performed a hierarchical clustering analysis to make clear the genetic differences and geographic variations among different individuals of the Tongyeong, Yeosu and Manjaedo population of *Pollicipes mitella*, respectively, collected in the southern sea by PCR analysis.

MATERIALS AND METHODS

Isolation of genomic DNA, primers and amplification conditions

Genomic DNAs were separated from 33 individuals of three turtle leg (*P. mitella*) populations of Tongyeong, Yeosu and Manjaedo located in the southern sea of Korea, as shown in Fig. 1. DNA extraction and/or purification were performed as described previously (Yoon, 2008). In order to achieve good results, DNA extraction should be performed with highest reagent (Bioneer Corp., Daejeon, Korea) and according to standard procedures with minor modification. After washings several times, 3 volumes of lysis buffer I (155 mM NH₄Cl; 10 mM KHCO₃; 1 mM EDTA) was added and mixed gently by inverting the tubes. The precipitates obtained were then centrifuged and resuspended in lysis buffer II (10 mM Tris-HCl, pH 8.0; 10 mM EDTA; 100 mM NaCl; 0.5% SDS), and 15 μ l of proteinase K solution (10 mg/ μ l) was added. After incubation, we added 300 μ l of 3 M NaCl, and tenderly pipetted for a few minutes. 600 μ l of chloroform was then added to the mixture and inverted (no phenol). Ice-cold 70% ethanol was added, and then the samples were centrifuged at 19,621 g for 5 minutes to extract the DNA from the lysates. The DNA pellets were then incubation-dried for more than 10 hours, maintained at -40°C until analysis. The purity and concentration of DNA was measured with the absorbance ratio by a spectrophotometer (Beckman DU 600 series, UK).

Seven primers, BION-45 (5'-CAAACGTCGG-3'), BION-52 (5'-GTGGAAGCGT -3'), BION-55 (5'-GTCACGGACG-3'),



Fig. 1. Genomic DNA samples isolated from three geographical *Pollicipes mitella* populations in Tongyeong, Yeosu and Manjaedo located in the southern sea of the Korean Peninsula.

BION-58 (5'-AGCCTGTGTC-3'), BION-67 (5'-GTAGACCCGT-3'), BION-76 (5'-GAGGCCCGTT-3') and BION-78 (5'-GGG GGTTAGG-3') were exposed to produce the unique loci to each population and number of shared loci by the three populations of turtle leg which could be evidently calculated. Amplified DNA analysis was performed using two Programmable DNA Thermal Cyclers (MJ Research Inc., Waltham, MA, USA). DNA amplification was performed in 25 μ l samples, which contained 10 ng of template DNA, 20 μ l of premix (Bioneer Corp., Daejeon, Korea), and 1 unit of primer. Amplification products were generated via electrophoresis on 1.4% agarose (Bioneer Corp., Daejeon, Korea) gel containing TBE (90 mM Tris, pH 8.5; 90 mM borate; 2.5 mM EDTA) with ethidium bromide. 100 bp ladder marker (Bioneer Corp., Daejeon, Korea) was used as a DNA molecular marker. The electrophoresed agarose gels were photographed by photoman direct copy system (PECA Products, Beloit, WI, USA) under

illumination. The degree of variability was calculated by use of the Dice coefficient (F), which is given by the formula: $F = 2 n_{ab} / (n_a + n_b)$, where n_{ab} is the number of bands shared between the samples a and b, n_a is the total number of bands for sample a and n_b is the total number of bands for sample b (Jeffreys & Morton, 1987; Yoon & Kim, 2004; Yoke-Kqueen & Radu, 2006). The relatedness between different individuals in the three population of turtle leg was generated according to the bandsharing values and similarity matrix. The hierarchical clustering tree was analyzed by the similarity matrices to generate a dendrogram using pc-package program Systat version 10 (SPSS Inc., Chicago, IL, USA).

RESULTS AND DISCUSSION

Genomic DNAs were isolated from the turtle leg (*P.*

mitella) populations of Tongyeong, Yeosu and Manjaedo located in the southern sea of the Korean peninsula. Authors assessed genetic variation among individuals of the Tongyeong, Yeosu and Manjaedo population of the turtle leg (*P. mitella*), respectively, collected in the southern sea by PCR analysis. DNA fragments acquired by seven decamer primers ranged in size from 50 to 1,600 bp in the turtle leg, as shown in Fig. 2. The bandsharing value between individuals no. 01 and no. 02 was 0.982 within the Tongyeong population, which was the highest value identified among the three populations (Table 1). The bandsharing value between individual's no. 13 of Yeosu population and no. 15 of Manjaedo population was 0.682, which was the lowest observed. All examined primers generated a total of 398 loci counted from the Yeosu population of the turtle leg, while a total of 385 loci were generated from the Manjaedo

Table 1. Similarity matrix including bandsharing values (BS) calculated using Nei and Li's index of the similarity of *Pollicipes mitella* from Tongyeong, Yeosu and Manjaedo of Korean peninsula, respectively

	BS from Tongyeong							BS from Yeosu							BS from Manjaedo						
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
1	-	0.982	0.928	0.921	0.945	0.817	0.962	0.868	0.785	0.826	0.778	0.822	0.695	0.814	0.790	0.802	0.800	0.806	0.777	0.790	0.785
2		-	0.948	0.915	0.940	0.94	0.893	0.878	0.77	0.825	0.802	0.834	0.703	0.825	0.801	0.813	0.811	0.79	0.789	0.804	0.784
3			-	0.886	0.927	0.904	0.945	0.831	0.727	0.775	0.818	0.834	0.686	0.800	0.759	0.79	0.786	0.753	0.742	0.796	0.742
4				-	0.940	0.915	0.919	0.778	0.715	0.778	0.763	0.775	0.700	0.799	0.769	0.771	0.771	0.737	0.72	0.747	0.779
5					-	0.958	0.961	0.808	0.735	0.792	0.801	0.799	0.744	0.822	0.764	0.774	0.775	0.756	0.742	0.748	0.789
6						-	0.961	0.839	0.768	0.812	0.816	0.804	0.737	0.839	0.769	0.777	0.777	0.771	0.756	0.752	0.797
7							-	0.856	0.745	0.818	0.818	0.808	0.725	0.843	0.780	0.808	0.807	0.787	0.787	0.782	0.802
8								-	0.846	0.825	0.826	0.88	0.717	0.830	0.780	0.809	0.795	0.813	0.789	0.802	0.768
9									-	0.871	0.810	0.842	0.720	0.825	0.787	0.704	0.788	0.815	0.802	0.795	0.781
10										-	0.785	0.796	0.701	0.841	0.755	0.814	0.774	0.826	0.788	0.791	0.844
11											-	0.897	0.73	0.858	0.826	0.813	0.764	0.807	0.805	0.794	0.775
12												-	0.719	0.884	0.852	0.874	0.828	0.814	0.829	0.850	0.806
13													-	0.723	0.680	0.721	0.693	0.691	0.713	0.686	0.668
14														-	0.898	0.849	0.817	0.84	0.823	0.842	0.831
15															-	0.872	0.841	0.817	0.816	0.874	0.816
16																-	0.921	0.895	0.912	0.916	0.852
17																	-	0.878	0.864	0.865	0.82
18																		-	0.845	0.858	0.767
19																			-	0.845	0.834
20																				-	0.783
21																					-




Fig. 2. Distribution of fragment sizes of Tongyeong, Yeosu and Manjaedo population of *Pollicipes mitella*. Solid lines: Tongyeong population. Dotted lines: Yeosu population. Thick dotted lines: Manjaedo population. The fragment numbers in each size interval have been computed from the pooled fragments obtained with all the primers. The higher fragment sizes (>1,200 bp) are much more observed in the Yeosu population.

Table 2. The number of average loci per lane by PCR analysis using 7 primers in *Pollicipes mitella* in Tongyeong, Yeosu and Manjaedo of Korea

Primer	Item	No. of average loci per lane		
		Tongyeong	Yeosu	Manjaedo
BION-45		7.1	8.7	9.1
		(50)	(61)	(64)
BION-52		6.4	4.6	3.9
		(45)	(32)	(27)
BION-55		5.9	5.6	6.6
		(41)	(39)	(46)
BION-58		7.3	8.4	8.4
		(51)	(59)	(59)
BION-67		9.7	9.1	8.7
		(68)	(64)	(61)
BION-76		10.3	11.6	10.7
		(72)	(81)	(75)
BION-78		10	8.9	9
		(70)	(62)	(63)
Total no.		397	398	395
Average no. per primer		56.7	56.9	56.4

The total number of loci, generated by 7 primers in *Pollicipes mitella* obtained from Tongyeong, Yeosu and Manjaedo, is shown in parentheses.

population, as illustrated in Table 2.

As regards average bandsharing value (BS) results, the turtle leg population from Tongyeong (0.929) exhibited higher bandsharing values than did turtle leg from Manjaedo (0.852), as illustrated in Table 4. This average

bandsharing value reported by our study is higher than the value reported for Spanish barbel species (0.71–0.81) (Callejas & Ochando 1998) and the average value between the two oyster populations (0.282 ± 0.008) (Kim et al., 2004). It seems to these authors that this is a result of a

Table 3. The number of unique loci to each population and number of shared loci by the three populations generated by PCR analysis using 7 decamer primers in Tongyeong, Yeosu and Manjaedo population of *Pollicipes mitella*, respectively

Primer\Population	Item	No. of unique loci to each population			No. of shared loci by the three populations
		Tongyeong	Yeosu	Manjaedo	Three populations (7 individuals per population)
BION-45		49	14	42	42
BION-52		35	21	14	21
BION-55		28	14	21	21
BION-58		42	42	28	84
BION-67		49	28	28	42
BION-76		70	56	63	147
BION-78		63	28	42	63
Total no.		336	203	238	420
Average no. per primer		48	29	34	60

Table 4. Multiple comparisons of average bandsharing values among Tongyeong, Yeosu and Manjaedo populations from three regions of Korean peninsula were generated according to the bandsharing values and similarity matrix

Population	Tongyeong	Yeosu	Manjaedo
Tongyeong	0.929	0.790	0.778
Yeosu	-	0.806	0.792
Manjaedo	-	-	0.852

high degree of inbreeding in narrow region for a long while. The number of unique loci to each population and number of shared loci by the three populations generated by PCR using 7 decamer primers in the turtle leg (*P. mitella*) population of Tongyeong, Yeosu and Manjaedo, as illustrated in Table 3. 336 numbers of unique loci to each population, with an average of 48 per primer, were observed in the Tongyeong population. 203 unique loci, with an average of 29 per primer, were identified in the Yeosu population. 238 unique loci, with an average of 34 per primer, were identified in the Manjaedo population. Especially, 420 numbers of shared loci by the three populations, with an average of 60 per primer, were observed in the three turtle leg (*P. mitella*) populations. Diagnostic markers that are found to be present in three populations of an eel-loach (*Pangio* sp.) are also considered to be species-specific markers, whereas the other bands were considered to be population-specific markers (Siti Azizah et al., 2005). The higher fragment sizes (>1,200 bp) are much more observed in the Yeosu population, as shown in Figs. 2 and 3.

Genetic distances among different individuals of the Tongyeong population of the turtle leg (lane 01–07), Yeosu population of the turtle leg (lane 08–14) and Manjaedo population of the turtle leg (lane 15–21), respectively, were generated using the CLASSIFICATION option in Systat version 10 according to the bandsharing values and similarity matrix, as shown in Fig. 4. The dendrogram, generated by seven trustworthy oligonucleotides primers, indicates three genetic clusters. Tongyeong population could be manifestly distinguished with the other two Yeosu and Manjaedo populations among three populations. The




Fig. 3. PCR-based electrophoretic profiles of individual *Pollicipes mitella*. Each lane shows DNA samples extracted from 21 individuals. DNA isolated from Tongyeong population (lane 1–7), Yeosu population (lane 8–14) and Manjaedo population of *P. mitella* (lane 15–21) were amplified by decamer primers BION-45 (A), BION-52 (B), BION-55 (C), BION-58 (D), BION-67 (E), BION-76 (F) and BION-78 (G). The PCR products were separated by 1.4% agarose gel electrophoresis and detected by ethidium bromide staining. 100 bp ladder marker was utilized as a DNA molecular size marker.




Fig. 4. Hierarchical dendrogram of genetic distances, obtained from three populations of *Pollicipes mitella*. The relatedness among different individuals in the *P. mitella* populations from Tongyeong, Yeosu and Manjaedo of the Korean peninsula were generated according to the bandsharing values and similarity matrix.

longest genetic distance (0.305) was found to exist between individuals' no. 02 of the Tongyeong population and no. 13 of the Yeosu population. In this study, genetic analysis has revealed a significant genetic distance among three turtle leg (*P. mitella*) populations. The shortest genetic distance displaying significant molecular differences was between individual's no. 01 and no. 02 of the Tongyeong population (0.018).

Three turtle leg (*P. mitella*) populations can be clearly distinguished, especially, by their morphological characters and PCR-based approach. In general, the population classification of turtle leg is based on morphological variations in body weight, body length, shell size, shell pattern, shell color, and shell length. It is assumed that differences in such characters reflect discrete origins or genetic characteristics (Chenyambuga et al., 2004). Above-mentioned, the prospective of oligonucleotides amplified polymorphic DNAs to identify diagnostic markers for species and population identification in teleosts (Callejas & Ochando, 1998; Yoon & Kim, 2004; Yoon, 2008) and in shellfish (McCormack et al., 2000; Kim et al., 2004; Park & Yoon, 2008) has also been well recognized.

ACKNOWLEDGEMENTS

The authors are grateful to the anonymous referees who assisted with thorough correction. Particular thanks also go to our laboratory colleagues, for their assistance in sample collection, DNA extraction/purification, electrophoresis, and their help with PCR techniques.

REFERENCES

- Callejas C, Ochando MD (1998) Identification of Spanish barbel species using the RAPD technique. *J Fish Biol* 53:208-215.
- Chenyambuga SW, Hanotte O, Hirbo J, Watts PC, Kemp SJ, Kifaro GC, Gwakisa PS, Petersen PH, Rege JEO (2004) Genetic characterization of indigenous goats of sub-Saharan Africa using microsatellite DNA markers. *Asian-Aust J Anim Sci* 17:445-452.
- Iyengar A, Piyapattanakorn S, Dtone DM, Heipel DA, Howell BR, Baynes SM, Maclean N (2000) Identification of microsatellite repeats in turbot (*Scophthalmus maximus*) and dover sole (*Solea solea*) using a RAPD-based technique: Characterization of microsatellite markers in dover sole. *Mar Biotechnol* 2:49-56.
- Jeffreys AJ, Morton DB (1987) DNA fingerprints of dogs and cats. *Animal Genetics* 18:1-15.
- Kim JY, Park CY, Yoon JM (2004) Genetic differences and DNA polymorphism in oyster (*Crassostrea* sp.) analysed by RAPD-PCR. *Korean J Genetics* 26:123-134.
- Klinbunga S, Ampayup P, Tassanakajon A, Jarayabhand P, Yoosukh W (2000) Development of species-specific markers of the tropical oyster (*Crassostrea belcheri*) in Thailand. *Mar Biotechnol* 2:476-484.
- McCormack GC, Powell R, Keegan B (2000) Comparative analysis of two populations of the brittle star *Amphiura filiformis* (Echinodermata: Ophiuroidea) with different life history strategies using RAPD markers. *Mar Biotechnol* 2:100-106.
- Park GS, Yoon JM (2008) Geographic variations between Jedo venus clam (*Protothaca jedoensis* Lischke) populations of Boryeong and Wonsan of Korea. *Korean J Malacol* 24:13-26.
- Siti Azizah MN, Ruzainah A, Patimah I (2005) Development of RAPD markers in the eel-loach (*Pangio* spp.) for genetic discrimination and monitoring of wild and cultured populations. *World Aquacul* 36:37-43.
- Welsh J, McClelland M (1990) Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res* 18:7213-7218.
- Welsh J, Petersen C, McClelland M (1991) Polymorphisms generated by arbitrarily primed PCR in the mouse: application to strain identification and genetic mapping. *Nucleic Acids Res* 19:303-306.
- Yoke-Kqueen C, Radu S (2006) Random amplified polymorphic DNA analysis of genetically modified organisms. *J Biotechnol* 127:161-166.

Yoon JM (2008) Variability in two species of Osmeridae (*Hypomesus nipponensis* and *Mallotus villosus*). Dev Reprod 12:151-158.

Yoon JM, Kim JY (2004) Genetic differences within and

between populations of Korean catfish (*S. asotus*) and bullhead (*P. fulvidraco*) analysed by RAPD-PCR. Asian-Aust J Anim Sci 17:1053-1061.