

# Partial Lipectomy of the Epididymal Fat Alters Expression of the Steroidogenic Enzymes in the Mouse Testis at Different Postnatal Ages

Yong-Seung Lee<sup>1\*</sup> and <sup>†</sup>Ki-Ho Lee<sup>2</sup>

<sup>1</sup>Department of Biological Science, Sungkyunkwan University, Suwon, Gyeonggi 16419, Korea

<sup>2</sup>Department of Biochemistry and Molecular Biology, College of Medicine, Eulji University, Daejeon 34824, Korea



\* Current address: Department of Microbiology and Immunology, University of Michigan Medical School, 1150 West Medical Center Drive, 5641 Medical Science II, Ann Arbor, MI, 48109-5620, USA.

Received: September 15, 2023

Revised: November 12, 2023

Accepted: November 30, 2023

## <sup>†</sup>Corresponding author

Ki-Ho Lee  
Department of Biochemistry and Molecular Biology, Eulji University, Daejeon 34824, Korea  
Tel: +82-42-259-1643  
Fax: +82-42-259-1649  
E-mail: kiholee@eulji.ac.kr

Copyright © 2023 The Korean Society of Developmental Biology.  
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ORCID

Yong-Seung Lee  
<https://orcid.org/0000-0002-8442-4024>  
Ki-Ho Lee  
<https://orcid.org/0000-0002-3495-5126>

## Conflict of interests

The authors declare no potential conflict of interest.

## Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2017R1D1A 1B03030209 and 2022R1F1A1076317).

## Abstract

The epididymal fat is a type of gonadal adipose tissue, which is localized closely to the testis. Even though it has been suggested that the epididymal fat is necessary for maintenance of spermatogenesis in the testis, the influence of epididymal fat on expression of testicular steroidogenic enzymes has not been examined. In the present research, expressional changes of steroidogenic enzymes in the mouse testis after 2 weeks of the surgical partial lipectomy of epididymal fat at different postnatal ages were determined by real-time polymerase chain reaction analysis. The transcript levels of all molecules at 2 months of postnatal age were significantly increased by the lipectomy of epididymal fat. However, the lipectomy at 5 months of postnatal age resulted in decreases of expression levels of all molecules examined in the testis. Except a reduced transcript level of hydroxysteroid 17-beta dehydrogenase 3, there were no significant changes of expression levels of other steroidogenic enzymes by the lipectomy at 8 months of postnatal age. At 12 months of postnatal age, the lipectomy caused a significant increase of transcript level of steroidogenic acute regulatory protein and a significant decrease of transcript level of hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1, without any expressional change of cytochrome P450 side chain cleavage, hydroxysteroid 17-beta dehydrogenase 3, and hydroxysteroid 17-beta dehydrogenase 3 in the testis. These findings suggest that the substances derived from epididymal fat could differentially influence on expression of steroidogenic enzymes in the testis during postnatal period.

**Keywords:** Epididymal fat, Testis, Steroidogenic enzymes, Real-time polymerase chain reaction (PCR), Lipectomy

## INTRODUCTION

The epididymal fat is a gonadal white adipose tissue and is found between the testis and head of epididymis (Niemała et al., 2008). Even though the epididymal fat-like structure in the mouse is observed from the birth, fully mature adipocyte of the epididymal fat is detectable from 14 days of postnatal age (Han et al., 2011). The epididymal fat is consisted of not only preadipocytes and mature

**Authors' contributions**

Conceptualization: Lee KH.  
Data curation: Lee YS.  
Formal analysis: Lee KH.  
Methodology: Lee KH.  
Investigation: Lee YS.  
Writing-review & editing: Lee YS, Lee KH.

**Ethics approval**

All procedures were performed in accordance with protocols approved by the Eulji University Animal Care and Use Committee (EUIACUC 20-04).

adipocytes but also various types of cells, including immune and endothelial cells (Cleary Margot et al., 1977). Like other types of white adipose tissue, the epididymal fat continuously grows and expands in mass and size during postnatal period, accompanying a changes of cellular compositions (Cleary Margot et al., 1977; Björntorp et al., 1979). The epididymal fat is divided into two regions, proximal and distal regions, depending on histochemical characteristics and the proximity to the testis (Tirard et al., 2007). But Altintas et al. (2011) have divided the epididymal fat into three parts relative to the location of spermatic blood vessels and have reported expression of several adipocytokines and the existence of different cellular compositions among these epididymal parts. These observations suggest the existence of regional differences within the epididymal fat, in the respect of behaviors and/or presumably functions.

Expression of adipogenic molecules in different epididymal fat parts during postnatal period has been examined (Lee, 2019; Lee & Kim, 2019). These researches have shown that expression patterns of adipogenic molecules differ between the epididymal fat regions according to postnatal age, even though the tendency of increased expression levels of adipogenic molecules has been observed with ages, in accordance with a steady growth of the epididymal fat during postnatal period (Lee, 2019; Lee & Kim, 2019). In addition, expression of several adipocytokines, including leptin, adiponectin, and resistin, in the epididymal fat at different postnatal ages has been evaluated (Lee & Kim, 2018; Lee, 2019; Lee & Kim, 2019). Similar with those of adipogenic molecules, expression of adipocytokines in the epididymal fat seems to be increased with postnatal age, in spite of distinct expression pattern of each adipocytokine during postnatal period (Lee & Kim, 2018; Lee, 2019; Lee & Kim, 2019). Thus, it is likely that differential expression of adipocytokines in the epididymal fat during postnatal period is accompanied with growth and development of the epididymal fat.

It is well recognized that adipose tissue is a highly active endocrine organ, which expresses and secretes various kinds of adipocytokines with endocrine functions (Kershaw & Flier, 2004). Many researches have determined the existence of receptors for adipocytokines in the testis, including leptin receptor (Tena-Sempere & Barreiro, 2002), adiponectin receptor (Ocón-Grove et al., 2008), and chemerin receptor (Li et al., 2014). In addition, the effects of adipocytokines on the testicular functions have been examined from many previous researches. Leptin induces decreases of steroidogenic acute regulatory protein (*Star*) and cytochrome P450 side chain cleavage (*Cyp11a1*) expression in the testis, thus resulting in reduction of testosterone production (Tena-Sempere & Barreiro, 2002). Chemerin also suppresses the production of testosterone by causing a decrease of 3 beta- and steroid delta-isomerase (*Hsd3b*) expression in the testis (Li et al., 2014). Moreover, a stimulatory effect of adiponectin on testosterone production in the testis by increasing expression of *Star* and *Cyp11a1* is found (Rak et al., 2017). These data clearly indicate that adipocytokines regulate testicular functions for testosterone production and spermatogenesis by controlling expression of steroidogenic enzymes. However, most of these effects of adipocytokines on the testis have been acquired from exogenous exposure to adipocytokines, manipulation of adipocytokine and its receptor genes, and/or induced clinical conditions, such as obesity. It is fairly difficult to find previous researches dealing with an innate influence of epididymal fat on the testicular functions. Chu et al. (2010) have demonstrated that surgical removal of the epididymal fat (EPFX) in the young adult hamster completely abolishes the spermatogenesis but not affects serum testosterone concentration. However, this research has not examined whether the EPFX gives an influence on expression of steroidogenic enzymes in the testis.

Thus the present research was aimed to find out the effect of partial EPFX on expression of testicular steroidogenic enzymes of mouse during postnatal period. Using a quantitative real-time polymerase chain reaction (PCR) analysis, the changes of expression patterns of steroidogenic enzymes in the testis by partial EPFX were evaluated and compared at different postnatal ages.

## MATERIALS AND METHODS

### 1. Experimental animals and surgical procedures of epididymal fat lipectomy

A total of 29 male C57BL/6 male mice at 7 weeks of postnatal age were obtained from Samtako (Osan, Korea). Each mouse was separately housed from others during a whole experimental period. The animals were permitted to access to food and water *ad libitum* and kept in controlled environments. Once reaching at 2 months of postnatal age, the animals were divided into four experimental age groups, a young adult group (2 months), two mature adults (5 and 8 months), and an elderly adult (12 months). Each experimental group was further divided into two sub-experimental groups, sham and epididymal fat lipectomy (EPFX) groups, including sham (n=4) and EPFX (n=4) at 2 months, sham (n=3) and EPFX (n=3) at 5 months, sham (n=3) and EPFX (n=4) at 8 months, and sham (n=4) and EPFX (n=4) at 12 months of postnatal age.

The surgical EPFX was performed to the animals at proper postnatal ages. The mouse was anesthetized with ether, and a small incision on each side of scrotum was made with a pair of fine scissors and forceps. The reproductive tract, including testis, epididymis, and epididymal fat, was exposed to the outside through the incision. The careful EPEX was performed to minimize interruption of the blood supply to the testis, and most of epididymal fat taken by EPEX was the proximal epididymal fat part. The equal surgical practice was carried out to another side of the male reproductive tracts. The scrotum incision was closed with instant glue (Aron Alpha Instant glue, Toagosei, Tokyo, Japan), and the careful observation was paid for assured recovery from the anesthetization and surgical actions. When the full recovery was confirmed, the mouse was placed back to its own cage and kept eye on any abnormal behaviors and casualty. Same surgical procedure was applied to the mouse in sham group, except EPFX.

### 2. Collection of the testis, isolation of total RNA from the testis, and generation of complementary DNA

The testis from each experimental mouse was collected after two weeks of the EPFX. The animal was euthanized by quick cervical dislocation, and the reproductive tract was isolated through an abdominal incision. In cold phosphate-buffered saline solution, the testis was rapidly separated from other parts of reproductive tract and quickly placed in liquid nitrogen. The testis was kept in  $-80^{\circ}\text{C}$  freezer for further use.

Isolation of total RNA from the frozen testis was carried out with Trizol reagent (Molecular Research Center, Cincinnati, OH, USA). Chloroform and isopropanol were sequentially applied to homogenized testis in Trizol solution to isolate and precipitate total RNA. Air-dried total RNA was re-suspended in DEPC- $\text{dH}_2\text{O}$ , and quantitative measurement of isolated total RNA was performed with NanoDrop Lite spectrophotometer (Thermo Scientific, Massachusetts, MA, USA). The quality of total RNA was evaluated by 1.2% agarose gel electrophoresis.

Generation of complementary DNA (cDNA) strand from isolated total RNA was carried out with 1  $\mu\text{g}$  of total RNA by using iScript<sup>TM</sup> Reverse transcription Supermix for reverse transcription (RT)-qPCR (Bio-Rad Laboratories, Hercules, CA, USA). The final volume of RT reaction was adjusted with nuclease-free  $\text{dH}_2\text{O}$  to 20  $\mu\text{L}$ , and the RT reaction was performed at  $25^{\circ}\text{C}$  for 5 min,  $46^{\circ}\text{C}$  for 20 min, and  $95^{\circ}\text{C}$  for 1 min. The cDNA product was directly used for a quantitative real-time PCR.

### 3. Quantitative real-time polymerase chain reaction (PCR) and statistical analysis

One microliter of cDNA generated was mixed with 7  $\mu\text{L}$  of iQ<sup>TM</sup> SYBR<sup>®</sup> Green Supermix (Bio-Rad Laboratories), 10 pmol of each oligonucleotide primer, and nuclease-free  $\text{dH}_2\text{O}$  to adjust to a total volume of 25  $\mu\text{L}$ . The examined steroidogenic enzymes include steroidogenic

acute regulatory protein (*Star*), cytochrome P450 side chain cleavage (*Cyp11a1*), cytochrome P450, family 17, subfamily A, polypeptide 1 (*Cyp17a1*), hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1 (*Hsd3a1*), and hydroxysteroid 17-beta dehydrogenase 3 (*Hsd17b3*), and the primer information utilized for real-time PCR is shown in Table 1. Glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*) was used for an internal control molecule for PCR analysis. The PCR condition was as follows, a pre-denaturation step at 95 °C for 5 min, cycles of a denaturation step at 95 °C for 30 sec, an annealing step at  $T_m$  for 30 sec, and an extension step at 72 °C for 30 sec in PTC-200 Chromo 4 real-time system (Bio-Rad Laboratories). An additional extension step at 72 °C for 10 min was added at the end of each PCR. The sizes of PCR products were confirmed by 1.2% agarose gel electrophoresis.

Independently triplicated RT reactions and real-time PCRs were performed to obtain a mean for each experimental animal, and the transcript levels of steroidogenic enzymes were adjusted to that of *Gapdh* to acquire means and standard errors of an experimental group. The data were expressed in the relative ratios of transcript levels of steroidogenic enzymes against *Gapdh*. The Student *t*-test was employed to determine statistical differences at transcript levels of steroidogenic enzymes between sham and EPFX groups at each postnatal age. If  $p < 0.05$ , it was considered to be statistically significant between two groups.

## RESULTS


### 1. Expressional changes of *Star* and *Cyp11a1* in the mouse testis after 2 weeks of epididymal fat lipectomy at different postnatal ages

The transcript level of *Star* in the testis of the EPFX mouse at 2 months of postnatal age was significantly higher than that of sham mouse (Fig. 1A). However, a significant decrease of *Star* transcript level was detected in the testis of the EPFX mouse at 5 months of postnatal age, compared with that in sham mouse (Fig. 1A). There was no significant change of *Star* mRNA abundance in the testis after the EPFX at 8 months of postnatal age (Fig. 1A). Interestingly, the EPFX at 12 months of postnatal age resulted in a significant increase of the abundance of testicular *Star* mRNA, compared to the level of sham group (Fig. 1A).

**Table 1. Primer information for real-time PCR analysis**

Gene	Sequence (5' → 3')	$T_m$ (°C)	PCR product (bps)
<i>Star</i> (NM_011485.5)	(F) GCTGGAAGTCCCTCCAAGACT (R) GCCACCCCTTCAGGTCAATA	60	61
<i>Cyp11a1</i> (AF195119.1)	(F) CGATACTTCTCATGCGAG (R) CTTTCTCCAGGCATCTGAAC	52	126
<i>Cyp17a1</i> (NM_007809.3)	(F) GGATCCGATACTGACTACCA (R) GGCTGGAGAGAGATTCACTA	50	212
<i>Hsd3b1</i> (NM_008293.4)	(F) GGATCATCAAGATGTTGGTGC (R) TGGTTTCTGGTCGGAATACTT	55	78
<i>Hsd17b3</i> (NM_008291.3)	(F) ATGGCATCGGAAAGCCTAT (R) CTCTTCTGCAATGGTCTGTAGC	60	111
<i>Gapdh</i> (NM_008084.3)	(F) GGCCATGAGGTCCACCACCCTGTTGCTGTA (R) CCCCTGGCCAAGGTCATCCATGACAACCTT	60	513

PCR, polymerase chain reaction; *Star*, Steroidogenic acute regulatory protein; *Cyp11a1*, cytochrome P450 side chain cleavage; *Cyp17a1*, cytochrome P450, family 17, subfamily A, polypeptide 1; *Hsd3a1*, hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1; *Hsd17b3*, hydroxysteroid 17-beta dehydrogenase 3; and *Gapdh*: glyceraldehyde-3-phosphate dehydrogenase.



**Fig. 1. Expression of *Star* and *Cyp11a1* in the mouse testis after 2 weeks of epididymal fat lipectomy at different postnatal ages.** The epididymal fat of lipectomized animal group (EPFX) was partially removed at 2 months (2M), 5 months (5M), 8 months (8M), and 12 months (12M) of postnatal age, and the testis was collected after 2 weeks of the lipectomy. The reproductive tract of sham group (sham) was exposed through the incision on the scrotum, and the incision was carefully sealed with the instant glue without the lipectomy of the epididymal fat. The levels of *Star* (a) and *Cyp11a1* (b) transcripts were normalized to those of *Gapdh*. Different letters on bars indicate statistical differences between experimental groups at  $p < 0.05$ . *Star*, Steroidogenic acute regulatory protein; *Cyp11a1*, cytochrome P450 side chain cleavage, and *Gapdh*, glyceraldehyde-3-phosphate dehydrogenase.

The testicular expression of *Cyp11a1* in the mouse at 2 months of postnatal age was significantly increased by EPFX (Fig. 1B). The abundance of *Cyp11a1* transcript in the testis of EPFX mouse at 5 months of postnatal age was significantly lower than that of sham group (Fig. 1B). Like *Star*, the EPFX did not influence on the level of testicular *Cyp11a1* transcript at 8 months of postnatal age (Fig. 1B). In addition, expression of *Cyp11a1* in the testis of EPFX mouse at 12 months of postnatal age was not significantly changed, compared to that of sham group (Fig. 1B).

## 2. Expressional changes of *Cyp17a1* and *Hsd3b1* in the mouse testis after 2 weeks of epididymal fat lipectomy at different postnatal ages


The expression of *Cyp17a1* in the mouse testis after the EPFX at 2 months of postnatal age was significantly increased, compared to that in sham group (Fig. 2A). However, a transient decrease of *Cyp17a1* transcript level was detected in the testis of EPFX group at 5 months of postnatal age (Fig. 2A). A significant decrease of *Cyp17a1* mRNA abundance in the testis of EPFX group was also found at 8 months of postnatal age, while there was no significant change of *Cyp17a1* expression level in the testis of EPFX group at 12 months of postnatal age (Fig. 2A).

The EPFX at 2 months of postnatal age resulted in a significant increase of *Hsd3b1* transcript level in the testis, compared to that in the testis of sham group (Fig. 2B). The mRNA abundance of the testicular *Hsd3b1* in the testis of EPFX mouse at 5 months of postnatal age was significantly lower than that of sham group (Fig. 2B). At 8 months of postnatal age, the EPFX did not give a significant influence on the level of the testicular *Hsd3b1* transcript (Fig. 2B). A significant decrease of *Hsd3b1* transcript level in the testis of EPFX mouse, compared to that of sham group was also found at 12 months of postnatal age (Fig. 2B).

## 3. Expressional change of *Hsd17b3* in the mouse testis after 2 weeks of epididymal fat lipectomy at different postnatal ages

The expression level of *Hsd17b3* in the testis of EPFX mouse at 2 months of postnatal age was






**Fig. 2. Expression of *Cyp17a1* and *Hsd3a1* in the mouse testis after 2 weeks of epididymal fat lipectomy at different postnatal ages.** The epididymal fat of lipectomized animal group (EPFX) was partially removed at 2 months (2M), 5 months (5M), 8 months (8M), and 12 months (12M) of postnatal age, and the testis was collected after 2 weeks of the lipectomy. The reproductive tract of sham group (sham) was exposed through the incision on the scrotum, and the incision was carefully sealed with the instant glue without the lipectomy of the epididymal fat. The levels of *Cyp17a1* (a) and *Hsd3a1* (b) transcripts were normalized to those of *Gapdh*. Different letters on bars indicate statistical differences between experimental groups at  $p < 0.05$ . *Cyp17a1*, cytochrome P450, family 17, subfamily A, polypeptide 1; *Hsd3a1*, hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1, and *Gapdh*, glyceraldehyde-3-phosphate dehydrogenase.

significantly increased, compared to that of sham mouse, while the level of *Hsd17b3* transcript in the testis of EPFX mouse was significantly decreased at 5 months of postnatal age (Fig. 3). At 8 months of postnatal age, the EPFX did not cause the change of *Hsd17b3* expression level in the mouse testis, and a similar observation was detected at 12 months of postnatal age (Fig. 3)

## DISCUSSION

Partial removal of the epididymal fat by surgical procedure at different postnatal age resulted in differential expression of steroidogenic enzymes in the testis, and the findings are as follows: 1) expression of all steroidogenic enzymes evaluated is significantly increased after EPFX at 2 months of postnatal age; 2) the EPFX caused significant decreases of transcript levels of testicular steroidogenic enzymes by EPFX at 5 months of postnatal age; and 3) the effect of EPFX at 8 and 12 months of postnatal ages on expression of steroidogenic enzymes in the testis is not likely prominent, even though there are some steroidogenic enzymes displaying significant changes on their expression levels.

Expression of adipocytokines in the epididymal fat has been examined by our and other researches. Altintas et al. (2011) have shown the existence of various adipocytokine transcripts in the mouse epididymal fat, including leptin, adiponectin, tumor necrosis factor (TNF)- $\alpha$ , and interleukin (IL)-6 and 10. Expression of resistin in the epididymal fat is also reported from a previous research (Ye et al., 2013). Our earlier studies have also shown expression patterns of leptin, adiponectin, and resistin transcripts in the epididymal during postnatal period (Lee, 2019; Lee & Kim, 2019). Notably our previous researches have displayed that expression patterns of these adipocytokines in distal and proximal epididymal fat parts are somewhat different from each other, especially at 5 months and 12 months of postnatal ages (Lee, 2019; Lee & Kim, 2019). In addition, others have demonstrated secretion of adipocytokines from the epididymal fat. Secretion



**Fig. 3.** Expression of *Hsd17b3* in the mouse testis after 2 weeks of epididymal fat lipectomy at different postnatal ages. The epididymal fat of lipectomized animal group (EPFX) was partially removed at 2 months (2M), 5 months (5M), 8 months (8M), and 12 months (12M) of postnatal age, and the testis was collected after 2 weeks of the lipectomy. The reproductive tract of sham group (sham) was exposed through the incision on the scrotum, and the incision was carefully sealed with the instant glue without the lipectomy of the epididymal fat. The level of *Hsd17b3* transcript was normalized to that of *Gapdh*. Different letters on bars indicate statistical differences between experimental groups at  $p < 0.05$ . *Hsd17b3*, hydroxysteroid 17-beta dehydrogenase 3 and *Gapdh*, glyceraldehyde-3-phosphate dehydrogenase.

of leptin from the epididymal fat tissue culture has been reported, and the amount of leptin secretion is higher at young adult than at immature animal (Nazian, 2013). Resistin secretion from the epididymal fat has been confirmed by *ex vivo* epididymal fat transplant culture (Ye et al., 2013). Together, these findings suggest that the epididymal fat could act as an endocrine tissue which releases various types of cytokines that might play a regulatory role in development and function of the male reproductive tract, including testis and epididymis. However, a direct effect of the epididymal fat on the male reproductive tract has been rarely examined, and the present research would offer the data indicating the possible functional role of the epididymal fat-derived adipocytokines on expressional regulation of steroidogenic enzymes in the testis.

The regulatory effects of adipocytokines on the testicular steroidogenesis by modulating expression of steroidogenic enzymes have been investigated by several researchers. Inhibitory effects of leptin and chemerin on expression of *Star*, *Cyp11a1*, and *Hsd3b*, accompanied with reduction of testosterone, in the testis have been examined (Tena-Sempere & Barreiro, 2002; Li et al., 2014), which adiponectin stimulates expression of testicular *Star* and *Cyp11a1* to induce the production of testosterone (Rak et al., 2017). The stimulatory effect of resistin on testosterone production has been determined by Nogueiras et al. (2004). However, these researches have applied with artificial conditions, such as exogenous exposure to adipocytokines and/or *in vitro* culture methods, and thus the outcomes from these researches would not be directly relevant to the effect of epididymal fat-derived adipocytokines on the testis. The influence of innate epididymal fat on the testicular functions has been determined by surgical lipectomy of epididymal fat (Chu et al., 2010). The EPFX causes the arrest of normal spermatogenesis but does not impact on testosterone level in serum (Chu et al., 2010). But, Chu et al. (2010) have not provide any information about expression of testicular steroidogenic enzymes by the EPFX. Our present study has shown differential responses on expression of steroidogenic enzymes in the testis to EPFX at different postnatal ages. Especially, the expressional responsiveness of testicular steroidogenic enzymes to EPFX is more notable at 2 and 5 months than at 8 and 12 months of postnatal ages. These observations imply

that expression of steroidogenic enzymes in the testis at younger age would be more susceptible to epididymal fat-derived adipocytokines than at old age. Moreover, the opposite expressional changes of steroidogenic enzymes at 2 and 5 months of postnatal ages suggest the existence of distinct regulatory mechanisms for expression of these testicular enzymes at two different postnatal ages.

Even though the current research has provided an evidence indicating the direct influence of the epididymal fat to expression of testicular steroidogenic enzymes for the first time to our knowledge, there are several issues to be considered for estimate of the physiological effect of epididymal fat on the testicular functions. Surgical removal of entire epididymal fat is almost impossible due to the distribution of innervation and blood systems present within the epididymal fat. The removed epididymal fat by EPFX in the present research is mostly the proximal, not distal, epididymal fat part. Thus, it is reasonable to consider that the findings from current study would be the effect of proximal epididymal fat part. The treatment of leydig cells with conditioned media collected from *in vitro* culture of distal or proximal epididymal fat part would provide an answer for regional effects of epididymal fat to the testis. In addition, a distinct responsiveness on expression of testicular steroidogenic enzymes to EPFX at different postnatal age should be further considered. The variable expression patterns of adipocytokine' receptors in the testis during postnatal period could be a possible suggestion. Moreover, a physiological effect of individual adipocytokine on the testis should be distinguished from others. Expressional inhibition of specific adipocytokine gene in the epididymal fat would give informative data for a defined role of the adipocytokine on regulation of the testicular functions, including steroidogenesis.

In summary, the current research has presented that the EPFX induces distinct expression patterns of testicular steroidogenic enzymes at different postnatal ages. Additional researches are suggested to clarify detailed physiological impact of the epididymal fat on the testicular functions during postnatal period.

## REFERENCES

- Altintas MM, Rossetti MA, Nayer B, Puig A, Zagallo P, Ortega LM, Johnson KB, McNamara G, Reiser J, Mendez AJ, Nayer A (2011) Apoptosis, mastocytosis, and diminished adipocytokine gene expression accompany reduced epididymal fat mass in long-standing diet-induced obese mice. *Lipids Health Dis* 10:198.
- Björntorp P, Karlsson M, Gustafsson L, Smith U, Sjöström L, Cigolini M, Storck G, Pettersson P (1979) Quantitation of different cells in the epididymal fat pad of the rat. *J Lipid Res* 20:97-106.
- Chu Y, Huddleston GG, Clancy AN, Harris RBS, Bartness TJ (2010) Epididymal fat is necessary for spermatogenesis, but not testosterone production or copulatory behavior. *Endocrinology* 151:5669-5679.
- Cleary Margot P, Greenwood MRC, Brasel JA (1977) A multifactor analysis of growth in the rat epididymal fat pad. *J Nutr* 107:1969-1974.
- Han J, Lee JE, Jin J, Lim JS, Oh N, Kim K, Chang SI, Shibuya M, Kim H, Koh GY (2011) The spatiotemporal development of adipose tissue. *Development* 138:5027-5037.
- Kershaw EE, Flier JS (2004) Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab* 89:2548-2556.
- Lee KH, Kim NH (2018) Expressional patterns of adipocyte-associated molecules in the rat epididymal fat during postnatal development period. *Dev Reprod* 22:351-360.
- Lee KH, Kim NH (2019) Differential expression of adipocyte-related molecules in the distal epididymal fat of mouse during postnatal period. *Dev Reprod* 23:213-221.
- Lee KH (2019) Postnatal expressional patterns of adipose-associated molecules in the mouse



- proximal epididymal fat. *Dev Reprod* 23:313-322.
- Li L, Ma P, Huang C, Liu Y, Zhang Y, Gao C, Xiao T, Ren PG, Zabel BA, Zhang JV (2014) Expression of chemerin and its receptors in rat testes and its action on testosterone secretion. *J Endocrinol* 220:155-163.
- Nazian SJ (2013) Leptin secretion from the epididymal fat pad is increased by the sexual maturation of the male rat. *J Androl* 22:491-496.
- Niemälä S, Miettinen S, Sarkanen JR, Ashammakhi N (2008) Adipose tissue and adipocyte differentiation: Molecular and cellular aspects and tissue engineering applications. In: Ashammakhi N, Reis R, Chiellini F (eds), *Topics in Tissue Engineering*. Vol. 4. Biomaterials and Tissue Engineering Group, Porto, Portugal, pp 1-26.
- Nogueiras R, Luz Barreiro M, Caminos JE, Gaytán F, Suominen JS, Navarro VM, Casanueva FF, Aguilar E, Toppari J, Diéguez C, Tena-Sempere M. (2004) Novel expression of resistin in rat testis: Functional role and regulation by nutritional status and hormonal factors. *J Cell Sci* 117:3247-3257.
- Ocón-Grove OM, Krzysik-Walker SM, Maddineni SR, Hendricks III GL, Ramachandran R (2008) Adiponectin and its receptors are expressed in the chicken testis: Influence of sexual maturation on testicular *ADIPOR1* and *ADIPOR2* mRNA abundance. *Reproduction* 136:627-638.
- Rak A, Mellouk N, Froment P, Dupont J (2017) Adiponectin and resistin: Potential metabolic signals affecting hypothalamo-pituitary gonadal axis in females and males of different species. *Reproduction* 153:R215-R226.
- Tena-Sempere M, Barreiro ML (2002) Leptin in male reproduction: The testis paradigm. *Mol Cell Endocrinol* 188:9-13.
- Tirard J, Gout J, Lefrançois-Martinez AM, Martinez A, Begeot M, Naville D (2007) A novel inhibitory protein in adipose tissue, the aldo-keto reductase AKR1B7: Its role in adipogenesis. *Endocrinology* 148:1996-2005.
- Ye H, Zhang HJ, Xu A, Hoo RLC (2013) Resistin production from adipose tissue is decreased in db/db obese mice, and is reversed by rosiglitazone. *PLoS One* 8:e65543.